

COMPSCI 514 Spring 2026: Midterm 2 Review

1 Logistics

- The exam is closed book, no calculator or cheatsheets allowed.
- Held **in class Thursday 4/23**.
- Format will be similar to Midterm 1: TRUE/FALSE questions, numerical answer questions (both with explanations optional), along with two longer ‘homework style’ questions.

2 Studying Suggestions

- Focus primarily on practice questions. First from the past finals, then from this sheet/questions I have asked in class, then from the textbooks if needed.
- On the practice finals, ignore questions about optimization (e.g., convexity, Lipschitzness, gradients, gradient descent, etc.), which we have not covered yet. On the Fall 2023 final (which was cumulative), ignore questions from the Randomized Algorithms unit.
- Make sure you understand all the homework and quiz problem solutions.
- Review slides as needed to fill in gaps in knowledge, but don’t make this your primary means of studying.

3 Concepts to Study

Probability and Randomized Algorithms (First Half of Class)

- The exam will not specifically test this part of the class, but you should be able to apply foundational techniques. E.g., compute expectations, linearity of expectation, union bound, take the expectation of a random matrix or a random dot product, etc.

Low Distortion Embedding and JL Lemma

- Low-distortion embeddings definition.
- The Johnson-Lindenstrauss Lemma, and how to apply it.
- Do not need to be able to recreate the JL proof, but should understand the ideas behind it and be comfortable thinking about random vectors and matrices as we do in the proof.

Low-Rank Approximation and PCA

- Understand and apply important linear algebraic manipulations used. E.g.:
 - $y^T y = \|y\|_2^2$ and using this to split $\|x - y\|_2^2 = \|x\|_2^2 + \|y\|_2^2 - 2x^T y$.
 - $\text{tr}(\mathbf{A}\mathbf{A}^T) = \text{tr}(\mathbf{A}^T \mathbf{A}) = \|\mathbf{A}\|_F^2 = \sum_{i=1}^{\text{rank}(\mathbf{A})} \sigma_i(\mathbf{A})^2$.
 - For $\mathbf{V} \in \mathbb{R}^{d \times k}$ with orthonormal columns, $\mathbf{V}^T \mathbf{V} = \mathbf{I}$ and $\mathbf{V}\mathbf{V}^T$ is a projection matrix.
 - By Pythagorean theorem $\|\mathbf{X} - \mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X}\mathbf{V}\mathbf{V}^T\|_F^2$.
 - Definition of eigenvectors and values.
 - Courant-Fischer theorem and connection to low-rank approximation.
- Low-rank approximation as projection onto a k -dimensional subspace. How this projection gives a compressed representation of a data matrix \mathbf{X} .
- Dual view of low-rank approximation as finding k vectors that approximately span the rows (data points) and the columns (features). High level understanding of why a data matrix may be nearly low-rank.
- Finding the best low-rank approximation (i.e., the best orthonormal span $\mathbf{V} \in \mathbb{R}^{d \times k}$) of \mathbf{X} using the eigenvectors of $\mathbf{X}^T \mathbf{X}$. Do not need to have full derivation memorized, but it is worth working through. Understand high level takeaways.
- Measuring the quality of the optimal low-rank approximation by plotting the eigenvalues (the spectrum).
- Ability to recognize when a matrix will be low-rank or close to low-rank (e.g., given in image, be able to make an educated guess about what its spectrum looks like.)
- Singular value decomposition definition.
- Connection of SVD of \mathbf{X} to eigendecompositions of $\mathbf{X}^T \mathbf{X}$ and $\mathbf{X}\mathbf{X}^T$. Connection of singular values to eigenvalues of $\mathbf{X}^T \mathbf{X}$ and $\mathbf{X}\mathbf{X}^T$.
- Computing PCA/optimal low-rank approximation from the SVD. Connection of left and right singular vectors to the dual view of low-rank approximation as row and column approximation.
- Low-rank approximation of a similarity matrix and entity embedding.
- Iterative methods for SVD: power method and high level ideas of analysis. When does it converge fast, when does it converge slow (up to what is covered on Thursday April 16 – nothing from Tuesday April 21).

Spectral Methods for Graphs

- Adjacency matrix \mathbf{A} and Laplacian ($\mathbf{L} = \mathbf{D} - \mathbf{A}$) definitions.
- Interpretation of the Laplacian as measuring how smooth a vector (a function) is over nodes of the graph.
- Connection to spectral embedding for non-linear dimensionality reduction.
- Motivation behind using the second smallest eigenvector of the Laplacian to find a small but balanced cut. $\vec{x}^T \mathbf{L} \vec{x}$ as giving the size of a cut when $\vec{x} \in \{-1, 1\}^n$ is a cut indicator vector.

- Graph clustering for non-linearly separable data and for community detection.
- Stochastic block model definition, expected adjacency matrix, Laplacian, and eigenvectors. Why spectral clustering works for stochastic block model.
- Understand the stochastic block model proof.

4 Practice Questions

1. Linear Algebra and Low-Rank Approximation

1. Exercises 3.6, 3.7, 3.8, 3.10, 3.11 (here $|\vec{x}|$ denotes the Euclidean norm of \vec{x}), 3.12, 3.13, 3.15, 3.18, 3.20, 3.21, 3.22, 3.26, 7.16, 12.31, 12.33 *Foundations of Data Science*.

2. Linear algebra practice:

- (a) For any vector \vec{y} verify that $\|\vec{y}\|_2^2 = \langle \vec{y}, \vec{y} \rangle = \vec{y}^T \vec{y}$.
- (b) If $\mathbf{X} = \mathbf{AB}$, \mathbf{X} 's columns are spanned by the columns of \mathbf{A} and \mathbf{X} 's rows are spanned by the rows of \mathbf{B} . Check that you understand why. What about when $\mathbf{X} = \mathbf{ABC}$ for some matrices $\mathbf{A}, \mathbf{B}, \mathbf{C}$. If $\text{rank}(\mathbf{A}) = k$, prove that $\text{rank}(\mathbf{X}) \leq k$.
- (c) For $\mathbf{V} \in \mathbb{R}^{n \times k}$ with orthonormal columns and vector $x \in \mathbb{R}^n$, $\|\mathbf{V}^T x\|_2 = \|x\|_2$. Always? Sometimes? Never?
- (d) Letting $\mathbf{U}_k \in \mathbb{R}^{n \times k}$ have columns equal to the top k left singular vectors of \mathbf{X} and $\mathbf{V}_k \in \mathbb{R}^{d \times k}$ have columns equal to the top k right singular vectors of \mathbf{X} , $\mathbf{U}_k \mathbf{U}_k^T \mathbf{X} = \mathbf{X} \mathbf{V}_k \mathbf{V}_k^T$. Always? Sometimes? Never?
- (e) Show that for any matrix \mathbf{A} with SVD $\mathbf{U} \mathbf{\Sigma} \mathbf{V}^T$,

$$\|\mathbf{A}\|_F^2 = \text{tr}(\mathbf{A}^T \mathbf{A}) = \text{tr}(\mathbf{A} \mathbf{A}^T) = \|\mathbf{U} \mathbf{\Sigma}\|_F^2 = \|\mathbf{V} \mathbf{\Sigma}\|_F^2 = \sum_{i=1}^n \sigma_i(\mathbf{A})^2,$$

where $\sigma_i(\mathbf{A})^2$ is the i^{th} singular value of \mathbf{A} (the i^{th} diagonal entry of $\mathbf{\Sigma}$) squared.


- (f) Prove the matrix Pythagorean theorem: that if $\mathbf{V} \in \mathbb{R}^{d \times k}$ has orthonormal columns, then for any matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$, $\|\mathbf{X} - \mathbf{X} \mathbf{V} \mathbf{V}^T\|_F^2 = \|\mathbf{X}\|_F^2 - \|\mathbf{X} \mathbf{V} \mathbf{V}^T\|_F^2$.
 - (g) For any $\mathbf{V} \in \mathbb{R}^{d \times k}$ with orthonormal columns, $\mathbf{V} \mathbf{V}^T$ is the projection matrix onto the subspace spanned by the columns of \mathbf{V} (\mathbf{V} 's column span). We used this fact many times when discussing low-rank approximation. Show that $\mathbf{V} \mathbf{V}^T = (\mathbf{V} \mathbf{V}^T)(\mathbf{V} \mathbf{V}^T)$. Why does this property make intuitive sense if $\mathbf{V} \mathbf{V}^T$ is a projection?
 - (h) Let $\mathbf{X} \in \mathbb{R}^{n \times d}$ be a matrix with singular values $\sigma_1(\mathbf{X}), \dots, \sigma_d(\mathbf{X})$ and SVD $\mathbf{X} = \mathbf{U} \mathbf{\Sigma} \mathbf{V}^T$. What are the eigenvalues of $\mathbf{X}^T \mathbf{X} + \lambda \mathbf{I}$. What are the corresponding eigenvectors?
 - (i) Show that if $v \in \mathbb{R}^n$ is an eigenvector of $\mathbf{A} \in \mathbb{R}^{n \times n}$ with eigenvalue λ , then so is $c \cdot v$ for any scalar c .
 - (j) Show that for any symmetric $\mathbf{A} \in \mathbb{R}^{n \times n}$ with eigenvalues $\lambda_1(\mathbf{A}), \dots, \lambda_n(\mathbf{A})$, and integer p , the eigenvalues of \mathbf{A}^p are equal to $\lambda_1(\mathbf{A})^p, \dots, \lambda_n(\mathbf{A})^p$.
3. Let $\mathbf{X} \in \mathbb{R}^{n \times d}$ have SVD $\mathbf{X} = \mathbf{U} \mathbf{\Sigma} \mathbf{V}^T$ with singular values $\sigma_1(\mathbf{X}), \dots, \sigma_d(\mathbf{X})$.
- (a) What are the eigenvalues of the matrix $(\mathbf{X}^T \mathbf{X})^2 + (\mathbf{X}^T \mathbf{X})^3$? What are its eigenvectors? How about the matrix $(\mathbf{X} \mathbf{X}^T)^2 + (\mathbf{X} \mathbf{X}^T)^3$?
 - (b) What is the runtime required to compute $[(\mathbf{X}^T \mathbf{X})^2 + (\mathbf{X}^T \mathbf{X})^3] \vec{v}$ for any $\vec{v} \in \mathbb{R}^d$.
 - (c) Name one method discussed in class which relies on efficiently applying a polynomial in $\mathbf{X}^T \mathbf{X}$ to a vector (or more generally, applying a polynomial in a matrix $\mathbf{A} \in \mathbb{R}^{d \times d}$ to a vector).

4. What is one reason why you would want to compute a low-rank approximation of a matrix $\mathbf{X} \in \mathbb{R}^{n \times d}$?
5. $\mathbf{X} \in \mathbb{R}^{500 \times 50}$ contains 500 well-clustered data points as its rows. In particular, there are ten cluster centers $\vec{y}_1, \dots, \vec{y}_{10} \in \mathbb{R}^{50}$, such that each row \vec{x}_i lies within Euclidean distance at most 1 of a center. Give an *upper bound* on $\min_{\mathbf{B}: \text{rank}(\mathbf{B})=10} \|\mathbf{X} - \mathbf{B}\|_F^2$.
6. Consider two matrices $\mathbf{A} = \begin{bmatrix} 1.01 & 0 \\ 0 & 1 \end{bmatrix}$ or $\mathbf{B} = \begin{bmatrix} 1.1 & 0 \\ 0 & 1 \end{bmatrix}$.
 - (a) What are their eigenvalues and eigenvectors?
 - (b) On which matrix will power method converge more quickly?
7. Let $\mathbf{X} \in \mathbb{R}^{n \times 900}$ have random entries drawn independently in $[0, 1]$. Let $\mathbf{Y} \in \mathbb{R}^{n \times 900}$ have rows corresponding to 30×30 pixel grayscale images of handwritten digits. Let $\mathbf{Z} \in \mathbb{R}^{n \times 900}$ have rows corresponding to 30×30 pixel grayscale images of handwritten letters from the English alphabet. All entries of \mathbf{Y} and \mathbf{Z} are in $[0, 1]$.
 - (a) How do you expect $\sum_{i=11}^{900} \sigma_i(\mathbf{X})^2$, $\sum_{i=11}^{900} \sigma_i(\mathbf{Y})^2$, and $\sum_{i=11}^{900} \sigma_i(\mathbf{Z})^2$ to compare? Explain why in a few sentences.
 - (b) Plot a guess at what the spectrums of these three matrices might look like. Do not worry about the scale of the y axis.
8. **A bit challenging:** Prove that the k^{th} singular value of any matrix \mathbf{A} is bounded as $\sigma_k(\mathbf{A})^2 \leq \frac{\|\mathbf{A}\|_F^2}{k}$. **Hint:** Use that $\|\mathbf{A}\|_F^2 = \sum_{i=1}^r \sigma_i(\mathbf{A})^2$. Use this to prove that for any integer $k > 0$ there exists \mathbf{B} with $\text{rank}(\mathbf{B}) = k$ such that $\|\mathbf{A} - \mathbf{B}\|_2^2 \leq \frac{\|\mathbf{A}\|_F^2}{k+1}$. Here $\|\mathbf{A} - \mathbf{B}\|_2$ is the spectral norm, given by $\|\mathbf{A} - \mathbf{B}\|_2 = \max_{x: \|x\|_2=1} \|(\mathbf{A} - \mathbf{B})x\|_2$ or equivalently $\|\mathbf{A} - \mathbf{B}\|_2 = \sigma_1(\mathbf{A} - \mathbf{B})$.

2. Spectral Methods for Graphs

1. Consider a graph G with Laplacian matrix \mathbf{L} . Consider the problem: $x_* = \arg \min_{x: \|x\|=1} x^T \mathbf{L} x$. What is x_* ? What value of $x_*^T \mathbf{L} x_*$ does it achieve?
2. Let G be a d regular graph (i.e., all vertices have d neighbors).
 - (a) What are the eigenvalues of G 's Laplacian \mathbf{L} in terms of the eigenvalues $\lambda_1(\mathbf{A}), \dots, \lambda_d(\mathbf{A})$ of its adjacency matrix \mathbf{A} ?
 - (b) What are the largest eigenvalue and and eigenvector of G 's adjacency matrix \mathbf{A} ?
3. Let G be a fully connected graph (a complete graph), with self-loops. What are the eigenvalues of its corresponding adjacency matrix \mathbf{A} and Laplacian \mathbf{L} ? What if there are no self-loops?
4. What is the sum of eigenvalues of the adjacency matrix of an undirected graph with no self-loops? What is the sum of squared eigenvalues? **Challenge question:** What is the sum of cubed eigenvalues? **Hint:** The first answer is a fixed number. The second depends on the number of edges in the graph. The third depends on the number of triangles in the graph.
5. Consider the stochastic block model.
 - (a) Why is clustering with the second largest eigenvector of the expected adjacency matrix equivalent to clustering with the second smallest eigenvector of the expected Laplacian?

- (b) Are these two approaches identical when clustering using the actual rather than the expected matrices?
 - (c) Describe a natural variant of the stochastic block model where these two algorithms would not be equivalent even on the expected matrices.
6. Describe in a sentence or two the difference between finding a minimum cut and partitioning a graph with the second smallest Laplacian eigenvector.
7. Consider the datasets below. You must run standard k -means clustering on one and spectral clustering on the other. Which would you apply each method to? Why?



8. Consider the two stochastic block model graphs shown below.
- (a) Which do you expect to have the largest spectral gap $\sigma_1(\mathbf{A}) - \sigma_2(\mathbf{A})$? Why?
 - (b) Which do you expect to have the lowest second-smallest Laplacian eigenvalue?

