

# CMPSCI 250: Introduction to Computation

Lecture #10: Partial Orders  
David Mix Barrington  
25 September 2013

# Partial Orders

- Definition of a Partial Order
- Total Orders
- The Division Relation
- More Examples of Partial Orders
- Hasse Diagrams
- The Hasse Diagram Theorem
- Proving the Hasse Diagram Theorem

## Definition of a Partial Order


- A **partial order** is a particular kind of binary relation on a set. Remember that  $R$  is a **binary relation** on a set  $A$  if  $R \subseteq A \times A$ , that is, if  $R$  is a set of ordered pairs where both elements of every pair are from  $A$ .
- Last time we used quantifiers to define four particular properties that a binary relation on a set might have.
- A relation is a partial order if and only if it is reflexive, antisymmetric, and transitive.

# Properties of a Partial Order

- A relation  $R$  is **reflexive** if every element is related to itself -- in symbols,  $\forall x: R(x, x)$ .
- It is **antisymmetric** if the order of elements in a pair can never be reversed unless they are the same element -- in symbols,  $\forall x: \forall y: (R(x, y) \wedge R(y, x)) \rightarrow (x = y)$ .
- Finally,  $R$  is transitive if  $\forall x: \forall y: \forall z: (R(x, y) \wedge R(y, z)) \rightarrow R(x, z)$ . This says that a chain of pairs in the relation must be accompanied by a single pair whose elements are the start and end of the chain.


# Diagrams of Binary Relations

- If  $A$  is a finite set and  $R$  is a binary relation on  $A$ , we can draw  $R$  in a diagram called a graph. We make a dot for each element of  $A$ , and draw an arrow from the dot for  $x$  to the dot for  $y$  whenever  $R(x, y)$  is true. If  $R(x, x)$ , we draw a loop from the dot for  $x$  to itself.





# Seeing the Properties

- The properties are perhaps easier to see in one of these diagrams.
- A relation is reflexive if its diagram has a loop at every dot.
- It is symmetric if every arrow (except loops) has a matching opposite arrow.




# Seeing the Properties

- It is antisymmetric if there are never two arrows in opposite directions between two different nodes.
- It is transitive if for every path of arrows (a chain where the start of each arrow is the end of the previous one) there is a single arrow from the start of the chain to the end.



# Clicker Question #1

- Which property does the diagrammed relation not have?




- (a) reflexive
- (b) symmetric
- (c) antisymmetric
- (d) transitive



# Answer #1

- Which property does the diagrammed relation not have?



- (a) reflexive
- (b) symmetric
- (c) *antisymmetric*
- (d) transitive



# Total Orders

- When we studied **sorting** in CMPSCI 187, we assumed that the elements to be sorted came from a type with a defined comparison operation.
- Given any two elements in the set, we can determine which is “smaller” according to the definition. (In Java the type would have a `compareTo` method or have an associated `Comparator` object.)

# Total Orders

- The “smaller” relation is not normally reflexive, but the related “smaller or equal to” relation is.
- Both these relations are normally antisymmetric, unless it is possible for the comparison relations to have ties between different elements.
- And both relations are transitive, just as  $\leq$  is on numbers.

# Total Orders

- But ordered sets have an additional property called being total, which we write in symbols as  $\forall x: \forall y: R(x, y) \vee R(y, x)$ .
- In general a partial order need not have this property -- two distinct elements could be incomparable.
- For example, the equality relation  $E$ , defined by  $E(x, y) \leftrightarrow (x = y)$ , is reflexive, antisymmetric, and transitive, but any two distinct elements are incomparable.

# The Division Relation

- Here's another example of a partial order that is not total.
- Our base set will be the natural numbers  $\{0, 1, 2, 3, \dots\}$ , and we define the division relation  $D$  so that  $D(x, y)$  means “ $x$  divides into  $y$  without remainder”.
- In symbols,  $D(x, y)$  means  $\exists z: x \cdot z = y$ . (Here we use the dot operator  $\cdot$  for multiplication.)

# The Division Relation

- Any natural divides 0, but 0 divides only itself.  $D(1, y)$  is always true.  $D(2, y)$  is true for even  $y$ 's (including 0) but not for odd  $y$ 's.  $D(100, x)$  is true if and only if the decimal for  $x$  ends in at least two 0's.
- In discussion next week we'll look at some tricks to determine whether  $D(k, y)$  is true for some particular small values of  $k$ .

## Division is a Partial Order

- It's easy to prove that  $D$  is a partial order.
- $D(x, x)$  is always true because we can take  $z$  to be  $1$  and  $x \cdot 1 = x$ .
- If  $D(x, y)$  and  $D(y, x)$  are both true,  $x$  must equal  $y$  because  $D(x, y)$  implies that  $x \leq y$ .
- And if  $D(x, y)$  and  $D(y, z)$ , then there exist naturals  $u$  and  $v$  such that  $x \cdot u = y$  and  $y \cdot v = z$ , and then we see that  $x \cdot (u \cdot v) = z$ .

## More Partial Order Examples

- There are several easily defined partial orders on strings.
- We say that  $u$  is a **prefix** of  $v$  if  $\exists w: uw = v$ . (Here we write concatenation as algebraic multiplication.) We say  $u$  is a **suffix** of  $v$  if  $\exists w: wu = v$ . And  $u$  is a **substring** of  $v$  if  $\exists w: \exists z: wuz = v$ .
- It's easy to check that each of these relations is reflexive, antisymmetric, and transitive.



## Clicker Question #2

- Let  $\Sigma$  be the alphabet  $\{a, b\}$  and consider the prefix, suffix, and substring relations on  $\Sigma^*$ . Which of these statements is *false*?
- (a)  $ab$  is a prefix of  $aba$  and  $aa$  is a substring of  $aba$
- (b)  $\lambda$  is a suffix of  $aba$  and  $\lambda$  is a substring of  $aba$
- (c)  $a$  is a suffix of  $aba$  and  $ba$  is a substring of  $aba$
- (d)  $aba$  is a prefix of  $aba$  and  $aba$  is a suffix of  $aba$

## Answer #2


- Let  $\Sigma$  be the alphabet  $\{a, b\}$  and consider the prefix, suffix, and substring relations on  $\Sigma^*$ . Which of these statements is *false*?
- (a) *ab is a prefix of aba and aa is a substring of aba*
- (b)  $\lambda$  is a suffix of aba and  $\lambda$  is a substring of aba
- (c) a is a suffix of aba and ba is a substring of aba
- (d) aba is a prefix of aba and aba is a suffix of aba

## More Partial Order Examples

- **Inclusion** on sets is another partial order, as  $X \subseteq X$ ,  $X \subseteq Y$  and  $Y \subseteq X$  imply  $X = Y$ , and  $X \subseteq Y$  and  $Y \subseteq Z$  imply  $X \subseteq Z$ .
- The **subclass** relation on Java classes is a partial order, since every class is a subclass of itself, two different classes can never each be subclasses of the other, and a subclass of a subclass is a subclass.


# More Partial Order Examples

- We represent this relation by an object hierarchy diagram in the form of a **tree**.
- One class is a subclass of another if we can trace a path of extends relationships in the diagram from the subclass up to the superclass.



# Hasse Diagrams


- We make a Hasse diagram by making a dot for each element of the set, and making lines so that  $R(x, y)$  is true if and only if there is a path from  $x$  up to  $y$ .
- (Relative position of points in a graph usually doesn't matter, but here it does.)



## Clicker Question #3

- To the left is a Hasse diagram for the division relation on part of  $\mathbf{N}$ . Which number could go in the place of  $x$ ?


- (a) 4
- (b) 6
- (c) 7
- (d) 9



# Answer #3


- To the left is a Hasse diagram for the division relation on part of  $\mathbf{N}$ . Which number could go in the place of  $x$ ?

- (a) 4
- (b) 6
- (c) 7
- (d) 9



# Hasse Diagram

- Starting from the graph of a partial order, we make a Hasse diagram as follows.
- We first delete the loops.
- We then position the nodes so the all arrows go upward.
- Finally we delete arrows that are implied by transitivity from other arrows.



Inclusion on Sets  
(wikipedia.org)



# The Hasse Diagram Theorem

- A Hasse diagram is a convenient way to represent a partial order if we can make one.
- But if I am just given  $R$  and told that it is a partial order, can I always make a Hasse diagram for it?
- The potential problem comes with the rule that the points must be arranged so that every arrow goes upward.

# The Hasse Diagram Theorem

- The **Hasse Diagram Theorem** says that any finite partial order is the “path-below” relation of some Hasse diagram, and the “path-below” relation of any Hasse diagram is a partial order.
- The second of these two statements is easy to prove -- we just have to check that the path-below relation is reflexive, antisymmetric, and transitive.
- We'll sort of prove the first statement here.

# Proving the Theorem

- Given the relation  $R$ , when do we want an arrow from  $x$  up to  $y$ ?
- There should be an arrow if  $R(x, y)$  is true and  $\neg\exists z: (x \neq z) \wedge (z \neq y) \wedge R(x, z) \wedge R(z, y)$ . (That  $z$  would make an  $x$ - $y$  arrow redundant.)
- To start drawing the diagram, we need an element that we can safely put at the bottom, because it has no arrows into it.

# Proving the Theorem

- An element  $x$  is called **minimal** for  $R$  if  $\forall y$ :  
 $R(y, x) \rightarrow (x = y)$ .
- A finite partial order must have at least one minimal element, because we can start somewhere and keep taking smaller elements until none exist.
- This process can't lead to a **cycle** because  $R$  is antisymmetric.

# Proving the Theorem

- We build the diagram recursively by finding a minimal element, making a Hasse diagram for the set without that element, and then putting the minimal element back at the bottom, with the arrows given by the rule above.
- To finish the proof, we have to make sure that the path-below relation of this diagram we've constructed is really  $R$ .