

# COMPSCI 514: Algorithms for Data Science

---

Cameron Musco

University of Massachusetts Amherst. Spring 2026.

Lecture 21

- Problem Set 4 is posted and due next Friday 5/8 at 11:59pm. .
- After today you should be able to solve all the problems.
- Final exam Tuesday 5/12.

# Summary

## Last Class:

- Multivariable calculus review
- Introduction to gradient descent. Motivation as a greedy algorithm.
- Convex functions
- Lipschitz functions

## This Class:

- Lipschitz functions
- Analysis of gradient descent for convex Lipschitz functions
- Extension to projected gradient descent for **constrained optimization**.

# Gradient Descent Psuedocode



## Gradient Descent

- Choose some initialization  $\vec{\theta}^{(0)}$ .
- For  $i = 1, \dots, t$ 
  - $\vec{\theta}^{(i)} = \vec{\theta}^{(i-1)} - \eta \nabla f(\vec{\theta}^{(i-1)})$
- Return  $\vec{\theta}^{(t)}$ , as an approximate minimizer of  $f(\vec{\theta})$ .

Step size  $\eta$  is chosen ahead of time or adapted during the algorithm (details to come).

# When Does Gradient Descent Work?

$$\theta \in \mathbb{R} \quad \nabla f(\theta) \in \mathbb{R}$$




Gradient Descent Update:  $\vec{\theta}_{i+1} = \vec{\theta}_i - \eta \nabla f(\vec{\theta}_i)$

# Convexity

**Definition – Convex Function:** A function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is convex if and only if, for any  $\vec{\theta}_1, \vec{\theta}_2 \in \mathbb{R}^d$  and  $\lambda \in [0, 1]$ :


$$(1 - \lambda) \cdot f(\vec{\theta}_1) + \lambda \cdot f(\vec{\theta}_2) \geq f\left((1 - \lambda) \cdot \vec{\theta}_1 + \lambda \cdot \vec{\theta}_2\right)$$



# Convexity

**Corollary – Convex Function:** A function  $f: \mathbb{R}^d \rightarrow \mathbb{R}$  is convex if and only if, for any  $\vec{\theta}_1, \vec{\theta}_2 \in \mathbb{R}^d$  and  $\lambda \in [0, 1]$ :

$$f(\vec{\theta}_2) - f(\vec{\theta}_1) \geq \vec{\nabla}f(\vec{\theta}_1)^T (\vec{\theta}_2 - \vec{\theta}_1)$$



## Practice with Definitions


Let  $f(\theta) = h(\theta) + g(\theta)$  where  $h$  and  $g$  are convex. Is  $f$  also convex?

## Practice with Definitions

Let  $f(\theta) = h(\theta) + g(\theta)$  where  $h$  and  $g$  are convex. Is  $f$  also convex?

## A second assumption: Lipschitzness

$$\theta \in \mathbb{R} \quad \nabla f(\theta) \in \mathbb{R}$$



Gradient Descent Update:

$$\vec{\theta}_{i+1} = \vec{\theta}_i - \eta \nabla f(\vec{\theta}_i)$$

Need to assume that the function is **Lipschitz** (size of gradient is bounded): There is some  $G$  s.t.:

$$\forall \vec{\theta} : \quad \|\vec{\nabla} f(\vec{\theta})\|_2 \leq G \Leftrightarrow \forall \vec{\theta}_1, \vec{\theta}_2 : \quad |f(\vec{\theta}_1) - f(\vec{\theta}_2)| \leq G \cdot \|\vec{\theta}_1 - \vec{\theta}_2\|_2$$

# Well-Behaved Functions

**Definition – Convex Function:** A function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is convex if and only if, for any  $\vec{\theta}_1, \vec{\theta}_2 \in \mathbb{R}^d$  and  $\lambda \in [0, 1]$ :

$$(1 - \lambda) \cdot f(\vec{\theta}_1) + \lambda \cdot f(\vec{\theta}_2) \geq f\left((1 - \lambda) \cdot \vec{\theta}_1 + \lambda \cdot \vec{\theta}_2\right)$$

**Corollary – Convex Function:** A function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is convex if and only if, for any  $\vec{\theta}_1, \vec{\theta}_2 \in \mathbb{R}^d$  and  $\lambda \in [0, 1]$ :

$$f(\vec{\theta}_2) - f(\vec{\theta}_1) \geq \vec{\nabla}f(\vec{\theta}_1)^T (\vec{\theta}_2 - \vec{\theta}_1)$$

**Definition – Lipschitz Function:** A function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is  $G$ -Lipschitz if  $\|\vec{\nabla}f(\vec{\theta})\|_2 \leq G$  for all  $\vec{\theta}$ .

# GD Analysis – Convex Functions

Assume that:

- $f$  is convex.
- $f$  is  $G$ -Lipschitz.
- $\|\vec{\theta}_1 - \vec{\theta}_*\|_2 \leq R$  where  $\vec{\theta}_1$  is the initialization point.

## Gradient Descent

- Choose some initialization  $\vec{\theta}_1$  and set  $\eta = \frac{R}{G\sqrt{t}}$ .
- For  $i = 1, \dots, t - 1$ 
  - $\vec{\theta}_{i+1} = \vec{\theta}_i - \eta \vec{\nabla} f(\vec{\theta}_i)$
- Return  $\hat{\theta} = \arg \min_{\vec{\theta}_1, \dots, \vec{\theta}_t} f(\vec{\theta}_i)$ .

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

**Step 1:** For all  $i$ ,  $f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{\|\vec{\theta}_i - \vec{\theta}_*\|_2^2 - \|\vec{\theta}_{i+1} - \vec{\theta}_*\|_2^2}{2\eta} + \frac{\eta G^2}{2}$ . **Visually:**

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

**Step 1:** For all  $i$ ,  $f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{\|\vec{\theta}_i - \theta_*\|_2^2 - \|\vec{\theta}_{i+1} - \vec{\theta}_*\|_2^2}{2\eta} + \frac{\eta G^2}{2}$ . Formally:

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

**Step 1:** For all  $i$ ,  $f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{\|\vec{\theta}_i - \vec{\theta}_*\|_2^2 - \|\vec{\theta}_{i+1} - \vec{\theta}_*\|_2^2}{2\eta} + \frac{\eta G^2}{2}$ .

**Step 1.1:**  $\vec{\nabla} f(\vec{\theta}_i)^T (\vec{\theta}_i - \vec{\theta}_*) \leq \frac{\|\vec{\theta}_i - \vec{\theta}_*\|_2^2 - \|\vec{\theta}_{i+1} - \vec{\theta}_*\|_2^2}{2\eta} + \frac{\eta G^2}{2} \implies$  **Step 1 by convexity.**

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

**Step 1:** For all  $i$ ,  $f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{\|\vec{\theta}_i - \vec{\theta}_*\|_2^2 - \|\vec{\theta}_{i+1} - \vec{\theta}_*\|_2^2}{2\eta} + \frac{\eta G^2}{2} \implies$

**Step 2:**  $\frac{1}{t} \sum_{i=1}^t f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{R^2}{2\eta \cdot t} + \frac{\eta G^2}{2}.$

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

Step 2:  $\frac{1}{t} \sum_{i=1}^t f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{R^2}{2\eta \cdot t} + \frac{\eta G^2}{2}.$

**Theorem – GD on Convex Lipschitz Functions:** For convex  $G$ -Lipschitz function  $f$ , GD run with  $t \geq \frac{R^2 G^2}{\epsilon^2}$  iterations,  $\eta = \frac{R}{G\sqrt{t}}$ , and starting point within radius  $R$  of  $\vec{\theta}_*$ , outputs  $\hat{\theta}$  satisfying:

$$f(\hat{\theta}) \leq f(\vec{\theta}_*) + \epsilon.$$

Step 2:  $\frac{1}{t} \sum_{i=1}^t f(\vec{\theta}_i) - f(\vec{\theta}_*) \leq \frac{R^2}{2\eta \cdot t} + \frac{\eta G^2}{2}.$